

## NEUTRON STARS AND THE FERMIONIC CASIMIR EFFECT

PIOTR MAGIERSKI

*Institute of Physics, Warsaw University of Technology, ul. Koszykowa 75  
Warsaw, PL-00662, Poland*

AUREL BULGAC

*Department of Physics, University of Washington  
Seattle, WA 98195-1560, USA*

PAUL-HENRI HEENEN

*Service de Physique Nucléaire Théorique, U.L.B - C.P. 229  
Brussels, B 1050, Belgium*

Received (received date)

Revised (revised date)

The inner crust of neutron stars consists of nuclei of various shapes immersed in a neutron gas and stabilized by the Coulomb interaction in the form of a crystal lattice. The scattering of neutrons on nuclear inhomogeneities leads to the quantum correction to the total energy of the system. This correction resembles the Casimir energy and turns out to have a large influence on the structure of the crust.

The average density of a typical neutron star is by 3 – 4 times larger than the density inside an atomic nucleus. Even though, in the neutron star crust the nucleon density is relatively low and do not exceed  $0.1 \text{ fm}^{-3}$ . The width of this layer reaches almost 10% of the star radius and its structure is quite complex. The outer part of the crust consists of nuclei forming a lattice immersed in an electron gas which become relativistic already at about  $10^6 \text{ g cm}^{-3}$ . Deeper in the star, in the inner crust, due to high density and pressure, the last occupied neutron levels in nuclei are no longer bound, and a neutron gas is formed. The structure of this region can thus be viewed as an inhomogeneous neutron matter, where nuclei play the role of impurities. The electrons at these densities are ultrarelativistic particles and therefore the electron-nucleus correlations, which are responsible for the complications of electronic structure in atoms and solids become unimportant. Eventually, at the bottom of the crust, the uniform nuclear matter is formed.

The structure of the outer parts of neutron stars are important for understanding of several observational issues. Namely, one expects that such phenomena as: thermal X-ray emission from the stellar surface, X-ray burst sources, or the sudden speed-ups in the rotation rate of some neutron stars may be related to its crustal properties. It provides thus a motivation for theoretical studies, and indeed the in-

vestigation of the crust structure has been the subject of a considerable theoretical effort (see e.g. <sup>1,2,3,4,5</sup> and reference therein). In particular, it has been found that in the inner crust the interplay between the Coulomb and the surface energies leads to the appearance of various nuclear phases, characterized by different shapes. An agreement has been reached on the existence of five phases which are formed in the region where the nucleon density varies from 0.03 to 0.1 fm<sup>-3</sup>. Most theoretical calculations predict a chain of phase transitions as the density increases: spherical nuclei → rods (“spaghetti” phase) → slabs (“lasagna” phase) → tubes → bubbles → uniform matter. However in all to date approaches shell effects have been either completely neglected, as in the models based on a liquid drop formula or Thomas-Fermi models, or have been taken into account only for bound nucleons. Namely, the total energy density of the system can be expressed schematically in the form:

$$E = E_{vol} + E_{surf} + E_{Coul} + E_{shell}, \quad (1)$$

where the volume term  $E_{vol}$  describes bulk properties of the neutron-proton-electron (npe) matter,  $E_{surf}$  denotes the surface energy of nuclei, and  $E_{Coul}$  takes into account the Coulomb interaction between protons and electrons. The last term denotes the shell energy and was usually omitted in the previous approaches. It represents the quantum correction to the total energy and has two origins. The first one is associated with bound nucleons. Since the nuclei form a lattice the single-particle bound states form narrow bands. Their distribution is not a smooth function of energy and therefore gives rise to an energy correction which is termed shell energy. The change of the total energy can be computed quite accurately using the shell correction method, once the single-particle spectrum is known <sup>6,7</sup>. It was shown however that this correction play a marginal role and does not influence the phase transition pattern <sup>2</sup>. The second effect is related to unbound neutrons which may scatter on inhomogeneities and form resonant states. The situation is somewhat similar to the Casimir effect in quantum field theory and condensed matter (see e.g. Refs. <sup>8,9,10</sup> and references therein), where the fluctuation induced interaction leads to an additional energy correction to the total energy of a system. In the same way, the scattering of unbound neutrons leads to an effective interaction between nuclei immersed in the neutron environment. This effect can be termed the fermionic Casimir effect and has been studied for simple geometries in Refs. <sup>11,12,13,14,15</sup>.

The shell energy for unbound neutrons can be calculated from the formula:

$$E_{shell}^{out} = \int_{-\infty}^{\mu} d\varepsilon \varepsilon g(\varepsilon, \mathbf{l}) - \int_{-\infty}^{\mu_0} d\varepsilon \varepsilon g_0(\varepsilon, \mathbf{l}). \quad (2)$$

In the above equation  $g_0(\varepsilon, \mathbf{l})$  stands for the Thomas-Fermi or liquid drop density of states related to the neutron gas and  $g(\varepsilon, \mathbf{l})$  is the true quantum density of states in the presence of inhomogeneities. An ensemble of parameters  $\mathbf{l}$  describes these objects and their relative geometrical arrangement. The parameters:  $\mu$  and  $\mu_0$  are

determined from the requirement that the system has a given average density of unbound neutrons:

$$\rho_n^{out} = \int_{-\infty}^{\mu} d\varepsilon g(\varepsilon, 1) = \int_{-\infty}^{\mu_0} d\varepsilon g_0(\varepsilon, 1). \quad (3)$$

Since in infinite matter the presence of various inhomogeneities does not lead to the formation of discrete levels, the effects we shall consider here arise from the scattering states, which is in complete analogy with the procedure for computing the Casimir energy. The only difference comes from the fact that the integration in the formula (2) is performed up to the Fermi energy instead of infinity, as in the expression for the Casimir energy. This fact is responsible for an oscillatory behavior of the shell energy as a function of the distance between nuclei.

If the impurities were small compared to the neutron Fermi wavelength then the generated interaction would originate mainly from the s-wave scattering giving rise to the Ruderman-Kittel correction which is small compared to the Fermi energy of neutrons<sup>16,17</sup>. However the s-wave scattering limit is not valid in this case since the neutron Fermi momentum  $k_F^n$  is of the order of  $1 fm$  and thus  $k_F^n R > 1$  where  $R$  is a size of an inhomogeneity. Consequently the contribution of higher partial waves to the scattering process cannot be neglected, giving rise to much larger energy correction, as for large objects more of the incident wave will be reflected<sup>11,12,13,14,15</sup>.

The determination of the shell energy requires the calculation of the scattering matrix of the system<sup>14</sup>. It is in principle a difficult task for arbitrary shapes and arbitrary mutual geometrical arrangements of inhomogeneities. It is possible however to obtain an approximate analytic result under simplifying assumptions concerning the nuclear shape. Namely, let us consider here only spherical, cylindrical and planar nuclei. Moreover since the amplitude of the wave function in the semiclassical limit is proportional to the inverse square root of the local momentum, the single particle wave functions for the unbound states will have a small amplitude over the deep well and thus the probability to find an unbound neutron inside the nuclei is reduced as compared to the surrounding space.<sup>a</sup> Hence one can approximately replace the nuclear potential by an effective repulsive potential of roughly the same shape<sup>18</sup>. Then the shell energy can be easily determined in the semiclassical approximation based on the Gutzwiller trace formula. Namely, the leading shell energy contribution to the total energy, for two obstacles being either spherical, rod-like or slab-like nuclei located at a distance  $d$  reads<sup>19,20</sup>:

$$E_{shell}^{out} \approx \frac{\hbar^2 L^i R^{2-i}}{8m_n} \left( \frac{3}{\pi} \right)^{\frac{2+i}{6}} \frac{(\rho_n^{out})^{\frac{2+i}{6}}}{d^{\frac{6-i}{2}}} \cos \left( 2k_F^n d - i \frac{\pi}{4} \right), \quad (4)$$

where  $m_n$  is the neutron mass and  $i = 0, 1, 2$  for two spherical, cylindrical and planar obstacles, respectively (we assume that rods and slabs are parallel to each

<sup>a</sup> There are of course a number of “resonant” delocalized states, whose amplitudes behave in an opposite manner. However, the number of such “resonant” states is small and brings only small corrections.




Fig. 1. The total energy density (1) of the npe matter as a function of the proton quadrupole moment  $Q_p = Q_{20}^p$ , calculated in the HF approximation at the constant nuclear density  $\rho$ , and proton fraction  $Z/A$ , where  $Z$  and  $A$  are the proton and nucleon numbers per cell, respectively. The lattice constant is equal to 26 fm (left subfigure) and 20.8 fm (right subfigure). The stable nuclear configurations are indicated by arrows.

other). In the above equation  $L$  defines the length of the obstacle, and  $R$  is its radius (in the case of a slab it is defined as half of its width).

Clearly, the interaction induced by shell effects depends on the nuclear shape and is a sensitive function of both the neutron density and the geometry of the mutual arrangement of nuclei. The interaction between many nuclei of various shapes may look, at first glance, quite complicated since three-, four-, and other many-body terms will appear as a result of multiple neutron scattering on inhomogeneities. One can show however that many-body terms are quite small and give merely a small corrections to the dominant pairwise interaction<sup>14</sup>.

The magnitude of the quantum corrections associated with the expression (4) is small as compared to the liquid-drop terms in eq. (1). What is important, however, is that the energy differences between various nuclear phases in the crust are also very small (of the order of a few  $keV/fm^3$ ), since the liquid drop terms almost cancel. One can see it in the figure 1, where the energy density of npe matter has been plotted as a function of nuclear deformation. The minima visible in the figure represent the stable nuclear phases corresponding to different shapes or lattice geometries. The results which are presented were obtained through the minimiza-

tion of the total energy functional (1) for the npe matter. Namely, we solved the Hartree-Fock equations for a fixed density of the npe matter with contributions in the energy functional coming from the nuclei, the neutron gas, and the electrons. In such a calculation, the liquid drop and the shell energy parts of the energy are automatically and self-consistently included. Since we solve the problem in a cubic box with three symmetry planes, several lattice geometries can be generated like e.g. simple cubic crystal (scc), face centered crystal (fcc), or body centered crystal (bcc) (see Refs. <sup>19,21</sup> for details).

Since the liquid drop energy for different phases is almost the same thus the shell energy plays a crucial role for determining the most energetically favored structure. In particular, it can be shown that the relative energies of different phases fluctuate rapidly as a function of the total density and these oscillations can be attributed to the shell effects associated with unbound neutrons <sup>19,20,11</sup>. Consequently, we may conclude that the structure of the crust may be quite complicated since the shell effects associated with unbound neutrons may easily reverse the phase transition order predicted by the liquid drop based approaches. Moreover the number of phase transitions may increase since the same phase may appear for various density ranges. It is also likely that the system will favor distorted lattices, or lattices with defects which decrease the shell energy.

Hence our results suggest that the purely quantum effect related to the fluctuation induced interaction between nuclei, and which thus may be termed the fermionic Casimir effect, is responsible for determining the structure of the inner crust of neutron stars.

## Acknowledgements

This research was supported in part by the Polish Committee for Scientific Research (KBN) under Contract No. 5 P03B 014 21 and the Wallonie/Brussels-Poland integrated action program. Numerical calculations were performed at the Interdisciplinary Centre for Mathematical and Computational Modelling (ICM) at Warsaw University. Authors would like to thank Hubert Flocard for providing us with the numerical code solving the Poisson equation. We also appreciate many discussions with A. Wirzba, Y. Yu, S.A. Chin and H. Forbert.

## References

1. G. Baym, H.A. Bethe, C.J. Pethick, Nucl. Phys. **A175**, 225 (1971).
2. K. Oyamatsu, M. Yamada, Nucl. Phys. **A578**, 181 (1994).
3. C.J. Pethick and D.G. Ravenhall, Annu. Rev. Nucl. Part. Sci. **45**, 429 (1995).
4. F. Douchin, P. Haensel, J. Meyer, Nucl. Phys. **A 665**, 419 (2000).
5. F. Douchin, P. Haensel, Phys. Lett. **B485**, 107 (2000).
6. V.M. Strutinsky and A. G. Magner, Sov. J. Part. Nucl. Phys. **7**, 138 (1976).
7. V.M. Strutinsky, Sov. J. Nucl. Phys. **3**, 449 (1966); Nucl. Phys. **A 95**, 420 (1967); *ibid* **A 122**, 1 (1968); M. Brack *et al.*, Rev. Mod. Phys. **44**, 320 (1972).

8. H.B.G. Casimir, Proc. K. Ned. Akad. Wet. **51**, 793 (1948); V.M. Mostepanenko and N.N. Trunov, Sov. Phys. Usp. **31**, 965 (1988) and references therein.
9. M.E. Fisher and P.G. de Gennes, C.R. Acad. Sci. Ser. B **287**, 207 (1978); A. Hanke *et al.*, Phys. Rev. Lett. **81**, 1885 (1998) and references therein.
10. M. Kardar and R. Golestanian, Rev. Mod. Phys. **71**, 1233 (1999) and references therein.
11. A. Bulgac, P. Magierski, Nucl. Phys. **A683**, 695 (2001).
12. A. Bulgac, P. Magierski, Phys. Scripta **T90**, 150 (2001).
13. A. Bulgac, P. Magierski, Acta Phys. Pol. **B32**, 1099 (2001).
14. A. Bulgac, A. Wirzba, Phys. Rev. Lett. **87**, 120404 (2001).
15. P. Magierski, A. Bulgac, Acta Phys. Pol. **B32** 2713 (2001).
16. M.A. Ruderman, C. Kittel, Phys. Rev. **96**, 99 (1954).
17. A.L. Fetter and J.D. Walecka, *Quantum Theory of Many Particle Systems*, (McGraw-Hill, New York, 1971).
18. A. Bulgac, S.A. Chin, H.A. Forbert, P. Magierski, Y. Yu in Proc. of the Int. Workshop on *Collective excitations in Fermi and Bose systems*", eds. C.A. Bertulani, L.F. Canto and M.S. Hussein, pp. 44-61, World Scientific, Singapore (1999); nucl-th/9811028.
19. P. Magierski, P.-H. Heenen, to be published.
20. P. Magierski, A. Bulgac, P.-H. Heenen, to be published in the Proc. of the Workshop: *The Nuclear Many-Body Problem 2001*, NATO Science Series II - Mathematics, Physics and Chemistry, Kluwer Academic Publishers.
21. P. Bonche, H. Flocard, P.-H. Heenen, S.J. Krieger, M.S. Weiss, Nucl. Phys. **A443**, 39 (1985).